

A photograph of a paved road curving through a natural landscape. The road is the central focus, curving from the bottom left towards the top right. The surrounding area is filled with dense, dark green vegetation, including trees and bushes. The sky is dark, suggesting a night or dusk setting. The overall mood is somber and mysterious.

**DE OVERSETE LEVESTEDER
ALVORLIGT OVERSETE...**

MORTEN DD HANSEN, NATURHISTORISK MUSEUM AARHUS



St Bredlundvej
Central Denmark Region
Street View - Oct 2010





Contents lists available at ScienceDirect

Biological Conservation

journal homepage: www.elsevier.com/locate/biocon



Optimizing grassland management for flower-visiting insects in roadside verges

Jinze Noordijk*, Katrien Delille, André P. Schaffers, Karlè V. Sýkora

Wageningen University, Nature Conservation and Plant Ecology Group, P.O. Box 47, 6700 AA Wageningen, The Netherlands

ARTICLE INFO

Article history:

Received 30 December 2008

Received in revised form 31 March 2009

Accepted 15 April 2009

Available online 20 May 2009

Keywords:

Diversity

Flower-abundance

Semi-natural grassland

Insect conservation

Mowing

Roadside verge

Vegetation management

ABSTRACT

The decline of flower-visiting insects is a threat to ecological processes and to the services these insects provide. Roadside verges in the Netherlands span approximately 80,000 km and are often covered with semi-natural grasslands. As such, they also provide a suitable habitat for many insects, but this has received little attention so far. We investigated the effects of different management treatments on flower-visiting insects. We studied flower visitation in a 3 years old experimental set-up with five mowing treatments each replicated five times. Management types were: no management and mowing once or twice per year with and without the removal of hay, representing common forms of management and neglect. During an entire growing season, both flowers (number of species and inflorescences) as well as insects (total abundance and actual flower visits) were investigated. Mowing twice per year with removal of hay showed highest values for all measured variables and this effect persisted throughout the growing season. The early summer cut proved to be very important for insect feeding opportunities, due to the re-flowering of plants later in the growing season. Flower abundance showed high correlations with both plant species richness and the number of insect visits. Although overall, mowing twice a year

Långawten 📍

← Ans, Central Denmark Region

🕒 Street View - Oct 2009



📍 Långawten

Back to Map






Navigation and thumbnail area. Includes a map thumbnail, a 'Back to Map' button, and a row of image thumbnails:

- Rørdal kirke fra 1930, beliggend...
- Gasværksvej, Aalborg, North De...
- Aalborg, North Denmark Region
- Gasværksvej, Aalborg, North De...
- Limfjord

Central Denmark Region
Street View - Aug 2010



Navigation controls including a compass, zoom in (+) and zoom out (-) buttons, and a 'Hide imagery' dropdown menu.




21 📍
← Rønde, Central Denmark Region
Street View - May 2009



Back to Map 📍

15, Rønde, Central Denmark Re... 15, Rønde, Central Denmark Re... 15, Rønde, Central Denmark Re... 15, Rønde, Central Denmark Re... 15, Rønde, Central Denmark Re... 15, Rønde, Central Denmark Re...





Øresundsmotorvejen
Capital Region of Denmark
Street View - Jul 2014













